

SHOULDER INJURIES IN WATER SPORT ATHLETES

Rapid therapy with minimal adverse effects

Canoe polo subjects the shoulders and trunk to extreme stresses. Shoulder injuries lead to unwelcome loss of time from sports. A rapid therapy with minimal adverse effects can help damaged structures heal.



Picture: © image images / iStock

RALPH KÜRSCHNER MD AND JAN SCHILLING MD /
ORTHOZENTRUM HAFENCITY, HAMBURG

According to the BMA guide to sport injuries [1], shoulder injuries in water sport athletes rank in the top five most common injuries. These include shoulder joint injuries with and without dislocation, rotator cuff injuries, disorders of the tendons (tendinitis) and arthritis.

These clinical pictures may have or may acquire degenerative and inflammatory components. To avoid the indicated surgery, where applicable, but also as follow-up treatment after surgery, another treatment that can be used alongside and supplementary to the generally accepted indicated treatments is infiltration with various substances. Corticoids were and are popularly used for injection treatments. Both the corticoids and local anaesthetics are effective in treating acute pain but are also reputed to be potentially cytotoxic and to hinder healing. On the other hand, the supply of oxygen and nutrients to damaged structures and their metabolic and structural integrity must be rapidly restored. Are there any (biological) alternatives?

BLOOD CELL SECRETOME (BCS)

BCS (Orthokine) is a cell-free cocktail of all the factors quantitatively released from the blood cells during coagulation. These factors initiate regeneration/wound healing after wound closure. They are released under controlled conditions at 37°C using the medical device EOT II. Although the healing effect cannot be attributed to one single factor alone, the role of the various growth factors in BCS in tissue healing has been established (e.g. TGF, HGF, IGF, PDGF, FGF). Other BCS components to which an effect can be attributed are cytokines, lipid mediators and extracellular vesicles.

CASE STUDY SHOULDER INJURY

A female water polo player (29) presented at our practice with pain and loss of strength in her left shoulder. She usually trained 6 times a week and had already struggled painfully through her last matches. Clinically, ultrasound

showed well marked subacromial bursitis, which was apparently not caused by trauma, and a partial supraspinatus tendon tear in the left shoulder (see also Fig.1). She received conservative treatment with targeted physiotherapy and osteopathy to improve/optimize function/biomechanics over a period of two months with concomitant gentle loading and cooling treatment. Puncture of the subacromial bursa and 1x subacromial bursa infiltration with corticoids were also performed.

Four weeks after the corticoid infiltration, BCS was infiltrated four times on the supraspinatus tendon under ultrasound control (every 3-4 days, 2 mL per infiltration). After the first infiltration, the patient already reported reduced pain, after the third, freedom from pain at rest, and after the fourth, only minimal pain under loading. Additional concomitant measures included function analysis, function training and dry needling. Ultrasound follow-up after three months showed marked structural improvement in the supraspinatus tendon. The patient continues to report freedom from pain under full loading. We attach ultrasound images for information purposes (Figs. 2 and 3), being well aware that ultrasound images are not comparable to "initial MRI images".

CURRENTLY AVAILABLE STUDIES ON BCS

A study on supraspinatus tendinopathy published in 2018 shows that infiltration with BCS is markedly superior to a corticoid in the medium term [2]. A series of earlier studies show improved healing of tendons and muscles with BCS in both humans and animals

[3-7]. These studies show histologically well marked improvement in tissue structure and, where investigated, significantly improved clinical findings. According to the currently available data, BCS improves the local metabolic environment inter alia by normalising cell physiology and reducing local oxygen radical formation. This allows local stem cells and macrophages to be active in tissue regeneration [8,9]. This property is also beneficial for acute and chronic joint injuries [10,11]. Post-operative treatment with BCS is also possible and significantly improves the outcome post ACL reconstruction [11].




FIG. 1 Before treatment: MRI of the affected left shoulder. Clearly visible bursitis and joint effusion as well as damage to the supraspinatus tendon.

JAN SCHILLING MD



is a specialist in orthopaedics and trauma surgery. The sports orthopaediatrician founded and manages the interdisciplinary Orthozentrum Hafencity. He has been providing care for squad and professional athletes for over 20 years. Among other things, he has been team doctor of various teams, including the FC St. Pauli football team, the Hamburg Freezers ice-hockey team, the HSV handball team and the Hamburg Crocodiles national hockey team. He has been tournament doctor at tennis tournaments for many years (including the ATP Hamburg European Open) and is a collaborative partner of the Hamburg Athletics Association.

RALPH KÜRSCHNER MD



is Orthopaedic and Trauma Surgery Consultant at Orthozentrum Hafencity, with additional qualifications in sports medicine and manual therapy. He is the team doctor of the first HTHC women's hockey team (Federal German women's hockey league), Euro Hockey League tournament doctor, tournament doctor at the beach volleyball world championships in 2019, collaborative partner of up-and-coming athletes, providing medical support, and tournament doctor at the ATP Hamburg European Open.

CONCLUSION

BCS is effective in soothing the painful inflammatory process (as in the described case) and, unlike corticoids, initiates regeneration of the damaged tissue. As the maintenance of the structure and function of the rotator cuff is crucial to the health of the shoulder joint, local BCS infiltrations are available as a supplement to the indicated established treatments to achieve a rapid and reliable return to match fitness.

References

- [1] Hardy et al. editors. *The BMA guide to sports injuries*. London: Dorling Kindersley; 2010
- [2] Damjanov et al. PMID: 30167587
- [3] von Wehren et al. PMID: 30900032
- [4] Majewski et al. PMID: 19875360
- [5] Genç et al. DOI: 10.1016/j.aott.2018.01.005
- [6] Geburek et al. PMID: 26113022
- [7] Wright Carpenter et al. PMID: 15532000 und PMID: 15532001
- [8] Blázquez et al. PMID: 31316380
- [9] Shirokova et al. PMID: 31847701
- [10] Baltzer et al. PMID: 18674932
- [11] Darabos et al. PMID: 21360125



FIG. 2 Before treatment: ultrasound image of the affected left shoulder. Visible supraspinatus tendon tear.



FIG. 3 Follow-up after 3 months: ultrasound image of the affected left shoulder. Visible structural improvement in the supraspinatus tendon tear.